

Number of Responders Correctly identifying surveillance for colorectal cancer in various diseases

CORRECT RESPONSE	PRE	POS T		PRE	POS T		PRE	POS T		PRE-TEST				POST-TEST					PRE	POS T	
SCREENING EXAM	PGY 1 n=22	PGY 1 n=20	P	PGY 2 n=15	PGY 2 n=11	P	PGY 3 n=13	PGY 3 n=10	P	PGY 1 n=22	PGY 2 n=15	PGY 3 n=13	P	PGY 1 n=20	PGY 2 n=11	PGY3 n=10	P	Total n=50	Total n=41	P	
Annual Flexible sigmoidoscopy beginning at age 10 years in FAP	8 (36.3%)	11 (55.0%)	0.2 26	2 (13.3%)	8 (72.7%)	0.004	3 (23.0%)	6 (60.0%)	0.10 2	8 (36.3%)	2 (13.3%)	3 (23.0%)	0.02 2	11 (55.0%)	8 (72.7%)	6 (60.0%)	0.06 48	13 (26.0%)	25 (60.9%)	0.001	
Annual Colonoscopy in Lynch Syndrome at 25 Years	16 (72.7%)	14 (70.0%)	0.8 45	12 (80.0%)	10 (90.1%)	0.6 14	9 (69.2%)	8 (80.0%)	0.66 0	16 (72.7%)	12 (80.0%)	9 (69.2%)	0.08 4 6	14 (70.0%)	10 (90.1%)	8 (80.0%)	0.4 40	37 (74.0%)	32 (78.0%)	0.654	
Annual Colonoscopy 8 years after diagnosis of IBD	20 (90.1%)	17 (85.0%)	0.6 56	13 (86.7%)	10 (90.1%)	1	10 (76.9%)	8 (80.0%)	1	20 (90.1%)	13 (86.7%)	10 (76.9%)	0.05 1 5	17 (85.0%)	10 (90.1%)	8 (80.0%)	0.8 61	43 (86.0%)	35 (85.4%)	0.932	